

Pediatrician's Phone Number : \_\_\_\_\_  
 Poison Control Phone Number: 1800-222-1222

Acetaminophen is the active ingredient in Tylenol.

Used to reduce fever and relieve pain.

DO NOT give to a child under 2 months of age.

Acetaminophen can be given every 4-6 hours.

Weight	Infant & Children's Tylenol Suspension (160 mg/5ml)	Tylenol Chewable (80 mg)
6-11 lbs.	$\frac{1}{4}$ tsp. or 1.25 ml.	-----
12-17 lbs.	$\frac{1}{2}$ tsp. or 2.50 ml.	-----
18-23 lbs.	$\frac{3}{4}$ tsp. or 3.75 ml.	-----
24-35 lbs.	1 tsp. or 5 ml.	2 tabs
36-47 lbs.	1 $\frac{1}{2}$ tsp.	3 tabs
48-59 lbs.	2 tsp.	4 tabs
60-71 lbs.	2 $\frac{1}{2}$ tsp.	5 tabs
72-95 lbs.	3 tsp.	6 tabs

Ibuprofen is the active ingredient in Motrin and Advil.

Used to reduce fever, swelling, and pain.

DO NOT give to a child under 6 months of age.

Ibuprofen can be given every 6-8 hours.

Weight	Infant's Ibuprofen Drops (50mg/1.25ml)	Children's Ibuprofen Suspension (100 mg/5ml)	Children's Ibuprofen Chewables (50 mg)	Junior Strength Ibuprofen Chewables or Tabs (100 mg)
12-17 lbs. (Greater than 6 months only!)	1.25 ml.	$\frac{1}{2}$ tsp.	-----	-----
18-23 lbs.	1.87 ml.	$\frac{3}{4}$ tsp.	-----	-----
24-35 lbs.	2.5 ml.	1 tsp.	-----	-----
36-47 lbs.	-----	1 $\frac{1}{2}$ tsp.	3 tabs	-----
48-59 lbs.	-----	2 tsp.	4 tabs	2 tabs
60-71 lbs.	-----	2 $\frac{1}{2}$ tsp.	5 tabs	2 $\frac{1}{2}$ tabs
72-95 lbs.	-----	3 tsp.	6 tabs	3 tabs

Benadryl's active ingredient is diphenhydramine

Used to reduce hives and itchy rashes.

May cause drowsiness, dizziness.

Benadryl can be given every 6-8 hours.

Weight	Benadryl Elixir (12.5mg/5ml)	Benadryl Chewables (12.5 mg)	Benadryl Tablets (25 mg)
12-17 lbs.	2.5 ml	-----	-----
18-23 lbs.	3.75 ml	-----	-----
24-35 lbs.	5 ml	1 chewable	-----
36-47 lbs.	7.5 ml	1 $\frac{1}{2}$ chewables	-----
48-59 lbs.	10 ml	2 chewables	1 tablet